

Hadronic contributions to the anomalous magnetic moment of the muon

Tom Blum,^a Peter Boyle,^{b,c} Mattia Bruno,^d Norman Christ,^e Davide Giusti,^f
Vera Gülpers,^c Taku Izubuchi,^{b,g} Luchang Jin,^{a,g} Andreas Jüttner,^h
Christoph Lehner,^{f,b*} Aaron S. Meyer,^b J. Tobias Tsangⁱ
(RBC and UKQCD collaborations)

October 1, 2020

Affiliations

^a Physics Department, University of Connecticut, Storrs, CT 06269, USA

^b Physics Department, Brookhaven National Laboratory, Upton, NY 11973, USA

^c Higgs Centre for Theoretical Physics, The University of Edinburgh, EH9 3FD, UK

^d Theoretical Physics Department, CERN, 1211 Geneva 23, Switzerland

^e Physics Department, Columbia University, NY 10024, USA

^f Universität Regensburg, Fakultät für Physik, 93040 Regensburg, Germany

^g RIKEN BNL Research Center, Brookhaven National Laboratory, Upton, NY 11973, USA


^h Physics and Astronomy, University of Southampton, Southampton SO17 1BJ, UK

ⁱ CP3-Origins and IMADA, University of Southern Denmark, Campusvej 55, 5230 Odense M, Denmark

*christoph.lehner@ur.de


- ▶ Persisting $3 - 4\sigma$ tension between the standard model theory and BNL measurement for muon $g-2 \Rightarrow$ Possible hint for new physics
- ▶ FNAL E989 will improve experimental precision of 0.54 ppm by a factor of four in the coming years. First results very likely by the end of this year.
- ▶ Current experimental uncertainty balanced with theory \Rightarrow theory uncertainty needs to be improved
- ▶ Theory uncertainty is dominated by two non-perturbative QCD contributions, the hadronic vacuum polarization (HVP) and the hadronic light-by-light scattering (HLbL) with similar individual uncertainties.

Hadronic vacuum polarization:




Source: Theory Initiative Whitepaper [arXiv:2006.04822](https://arxiv.org/abs/2006.04822)


Isospin-symmetric limit:




QED corrections:



Strong isospin-breaking corrections:




- ▶ Need to calculate at 1-2 per-mille precision to match experimental precision after Fermilab E989. Current status: first results around 1%.
- ▶ Further error reduction by factor of 10 is a challenging task for the next 5-10 years.
- ▶ Many improvements for improved statistical and systematic control have already been developed, e.g., improved bounding method (RBC/UKQCD) for light-quark connected isospin symmetric contribution:



Future:

- ▶ Need statistics dominated error budgets!
- ▶ Simulations at finer lattice spacings a with $a^{-1} \approx 5$ GeV desired! See related computational Lol with algorithmic developments to accelerate this.
- ▶ Within RBC/UKQCD, simulations are already only at physical pion mass, so no chiral extrapolation needed. High-precision control of QED and SIB effects crucial.
- ▶ Combined R-ratio/lattice QCD approaches as proposed by RBC/UKQCD 2018 may play a crucial role to scrutinize and improve the estimates. Euclidean time windows also serve as tools for cross-checks.

Hadronic light-by-light scattering:



First ab-initio result this year (RBC/UKQCD
[PRL124\(2020\)132002](#)):

$$a_{\mu}^{\text{HLbL}} 10^{10} = 7.87(3.06)_{\text{stat}}(1.77)_{\text{sys}} \quad (1)$$

For this quantity it will suffice to reduce the uncertainty to $O(10\%)$ in the next 5-10 years.

Future:

- ▶ Our previous result used a finite-volume regulator for the photons, will focus on infinite-volume photons coupled to finite-volume quark correlators.
- ▶ This removes large power-law finite-volume errors but requires special attention to control statistical uncertainties originating from long-range contributions.
- ▶ For this we will re-use successful ideas of our HVP projects with regards to full low-mode averaging using a locally coherent Dirac eigenmodes representation.
- ▶ We aim to also calculate the long-distance part of the neutral pion-pole contribution separately to further reduce the statistical noise and finite-volume dependence.
- ▶ In the current calculation some sub-leading diagrams are only estimated to be small, we will verify this explicitly.
- ▶ We have previously shown that different infinite-volume regulators can be used to reduce systematic uncertainties. We will explore this further.

Summary of precision goals for next 5-10 years:

HVP: 1% \rightarrow 0.1%

HLbL: 44% \rightarrow 10%

What is required to succeed? We need continued access to large-scale computing resources in the US and dedicated support for the lattice QCD project.